

P. SRIVASTAVA, G. KUMAR

Department of Botany, Plants Genetics Laboratory,
University of Allahabad, Allahabad 211 002, INDIA

Email address of corresponding author- *prisri.aks1985@gmail.com

*Address of corresponding author: C/o Dwarka Prasad Srivastava
13/14 Karanpur (Near Prayag Railway Station), Allahabad 211 002 India.

EMS-INDUCED CYTOMICTIC VARIABILITY IN SAFFLOWER (*CARTHAMUS TINCTORIUS* L.)



Seeds of safflower (Carthamus tinctorius L.) were subjected to three treatment durations (3h, 5h and 7h) of 0.5 % Ethyl Methane Sulphonate (EMS). Microsporogenesis was carried out in the control as well as in the treated materials. EMS treated plants showed interesting feature of partial inter-meio-cyte chromatin migration through channel formation, beak formation or direct cell fusion. Another interesting feature noticed during the study was the fusion among tetrads due to wall dissolution. The phenomenon of cytomixis was recorded at nearly all the stages of microsporogenesis connecting from a few to several meiocytes. Other abnormalities such as lag-gards, precocious movement, bridge and non-disjunction of chromosomes were also recorded but in very low frequencies. The phenomenon of cytomixis increased along with the increase in treatment duration of EMS. Cells with these types of cytotoxic disturbances may probably result in uneven formation of gametes or zygote, heterogenous sized pollen grains or even loss of fertility in future.

© P. SRIVASTAVA, G. KUMAR, 2011

Introduction. Since the first discovery of cytomixis in *Crocus sativus* by Kornicke [1] it has been reported in a number of plant species [2–7], but reports on cytotoxic variations in safflower is meager due to pure staining ability, problem of chromosome stickiness and lack of research work on its cytogenetical aspects. The phenomenon of cytomixis is characterized by the migration of chromatin/chromosomes between the surrounding meiocytes through cytoplasmic channels or direct fusion among PMCs.

Although cytomixis has been reported in a number of plant species, its origin is still an enigma. Among the factors proposed to cause cytomixis are: 1) the influence of genes [8]; 2) abnormal formation of the cell wall during premeiotic divisions [9]; 3) action of chemical agents such as colchicine and methyl methane sulphonate [10] rotenone [11]; 4) changes in the biochemical process that involves microsporogenesis modifying the microenvironment of affected anthers [12]; 5) the effect of gamma radiation resulting in an imbalanced and sterile genetic system [13]; 6) the presence of a male-sterile mutant gene and its frequency altered by environmental factors [14], or environmental stress, and pollution [15].

The persistence of cytotoxic channels has significant effect on the meiotic process, its end products and fertility of the plant. This process seems to be of evolutionary significance, which is endorsed by many authors [16, 17] and is more frequent in species with unbalanced genomes such as haploids, aneuploids, hybrids [18], mutants [2] triploids [19] and apomicts [20]. It may lead to the production of aneuploid plants with certain morphological characteristics [21] or produce unreduced gametes as reported in several grass species including *Dactylis* [22] and *Aegilops* [23]. Unreduced gamete formation is of evolutionary importance leading to the production of plants with higher ploidy levels. Aneuploid pollen grains produced as a result of cytomixis may also be responsible for reduction in pollen fertility of the *Stipa* species [24].

Material and Methods. Seeds of *C. tinctorius* L. were obtained from the Genetics Division, Indian Agricultural Research Institute (IARI), New Delhi. Healthy seeds of the plant were pre-soaked in distilled water and then treated with ethyl methane sulphonate (EMS) at three treatment durations (3 h, 5 h and 7 h) of 0.5 % EMS. The treated seeds were washed in running tap water to remove the residual effect of the mutagen sticking

to the seed coat. One set of seeds was kept untreated to act as control for comparison. For meiotic studies, young flower buds were randomly selected and then fixed in freshly prepared Carnoy's fluid (alcohol: glacial acetic acid in a 3:1 ratio) for 24h and preserved in 70 % alcohol at 4 °C. The anthers were squashed in 2 % acetocarmine, and photographs were taken from freshly prepared slides using Nikon Image capturing system.

Results. The meiosis in control plants was absolutely normal with 12 bivalents at metaphase-I and 12:12 separation at anaphase-I without any evidence of chromosomal anomalies or cytotoxic variations. Cytomixis, the inter PMC (Pollen Mother Cell) transfer of chromatin material is observed only in the populations treated with EMS. Data regarding types of karyological abnormality, cytotoxic abnormality, total cytoplasmic abnormality and pollen fertility is provided in the Table. The frequency and intensity of cytotoxicity depended on the nature of connection between adjacent cells. Maximum amount of cytotoxicity is contributed to inter-chromatid migration through tube or beak formation (Fig. 2, 3) with a frequency of 1.95 % at 7 h treatment duration followed by cytotoxicity among PMCs either at same phase of division (Fig. 1, 4, 6) or at different phase (Fig. 5) being 0.85 %, then comes cytotoxicity through wall dissolution between pollen tetrads being 0.80 % (Fig. 8). Inter-meioocyte transfer of genetic material as micronuclei was also noted in considerable amount (Fig. 7). Total frequency of cytoplasmic connections was recorded to be 0.85 ± 0.66 %, 1.62 ± 0.53 % and 3.55 ± 0.72 % at 3, 5, 7 h treatment duration respectively as also depicted in the Fig. 9.

As regard the karyological abnormality percentage, abnormal meiotic course is characterized by the occurrence of chromatin stickiness predom-




Fig. 1. Migration of chromosome between same phase of three PMCs




Fig. 2. Cytotoxicity through Tube formation



Fig. 3. Cytotoxicity through Beak formation

EMS induced cytotoxic variations in *Carthamus tinctorius* L.

Treatment dose (h)	No. of PMCs Scored	Karyological abnormality, %	Cytoplasmic abnormality, %			Total cytoplasmic abnormality, %	Pollen fertility, %
			cy	icm	cwd		
Control	465	—	—	—	—	—	96.95
3h	451	8.33	0.38	0.25	0.22	0.85 ± 0.66	88.15
5h	463	17.65	0.32	0.95	0.35	1.62 ± 0.53	70.92
7h	492	32.16	0.85	1.95	0.80	3.55 ± 0.72	68.65

Abbreviations: cy—cytotoxicity between PMCs, icm—inter chromatid migration, cwd—cytotoxicity through wall dissolution.




Fig. 4. Cytomixis through cell wall dissolution between similar stages involving movement of extra chromosomal mass




Fig. 5. Cytomixis between more two PMCs of dissimilar stages




Fig. 6. Cytomixis between two PMCs of similar stages (metaphase-I)

inantly at MI and unequal separation of bivalents/chromosomes at AI/II, bridges and laggards at anaphases were also present in significant amount (Table). Consequent effect of these abnormalities on the microsporogenesis is evident from the occurrence of abnormal tetrads configurations and microsporocytes with micronuclei.

The test for pollen fertility showed fewer amounts of sterile pollen grains in control plants as




Fig. 7. Cytomixis involving elimination of nucleic material as micronuclei




Fig. 8. Direct fusion of tetrads due to cell wall dissolution

compared to treated sets. Pollen fertility recorded in control plants was found to be 96.65 % which decreased to 68.65 % at 7h treatment duration sets. No significant correlation was found between cytomixis and pollen viability.

Discussion. Cytomixis, the migration of chromatin material between and among the meiocytes, meristematic, tapetal, integumental, nucellar and ovary cells through cytomictic channels or direct adhesion is a well established phenomenon reported in large array of plants [5, 7, 25]. Cytomixis is a potential means to conserve the genetic heterozygosity of gametes [26] and additional means of phylogenetic evolution of karyotypes by reducing or increasing the basic series [27, 28] creation of aneuploids and polyploids [29]. This is a peculiar type of cytological anomaly reported in number of angiosperm species [30, 31]. According to Heslop-Harrison [32] and Risueno et al. [33] the role of cytoplasmic channels is related to transport of nutrients between meiocytes. Although the phenomenon of cytomixis is reported in a number of plant species but the




Fig. 9. Percentage of cytotoxicity at three different time durations of 0.5 % EMS

chromosome transfer in true sense is rare. Kamra [9] working on PMC of *Hordeum* believed that no amount of defective squashing or application of pressure could produce such small protusions so close to one another or to form PMCs with extra fragments or increase in the number of bivalents in them especially at metaphase. In recent years there is accumulating evidence that cytotoxicity is a normal, genetically controlled phenomenon influenced by physiological and environmental factors [7, 35, 36] rather than being due to fortuitous causes such as fixation, mechanical injuries or pathological anomaly. Cytotoxicity as a cause of inducing meiotic irregularities and formation of hypo- and hyperploid PMCs has also been reported in several plants [29, 30, 32, 37]. The study revealed that all the stages of meiosis were susceptible to cytotoxicity as contrary to the general belief that early stages are more favourable [38].

The possible implications of cytotoxicity are 1. either it may give rise to pollen sterility due to degeneration of cells with no or very little genetic material and 2. most of the cells completing all meiotic stages, would be genetically imbalanced because of less or more number of chromosomes than normal ones [39].

All of the meiotic abnormalities found in the EMS treated set of safflower analyzed revealed that there was no correlation between cytotoxicity and decrease in pollen fertility percentage but type of meiotic anomalies had direct correlation with pollen fertility. Fertility depends on the efficiency of the meiotic process. Increase in meiotic aberrations caused decrease in pollen fertility percentage which was in concordance with studies conducted by various workers on *Allium*, *Pilocarpus* [40–43].

I would like to extend my thanks to my seniors

who supported my work with their valuable suggestions and guidance. I am also grateful to Mrs. Ritambhara Shukla for critical reading of the manuscript.

P. Srivastava, G. Kumar

ЦИТОМИКТИЧЕСКАЯ ВАРИАБЕЛЬНОСТЬ
У САФЛОРА (*CARTHAMUS TINCTORIUS* L.),
ИНДУЦИРОВАННАЯ ЭМС

Семена сафлора (*Carthamus tinctorius* L.) обрабатывали 0.5%-ным этилметансульфонатом (ЭМС) в одном из трех режимов – 3, 5 и 7 ч. Микроспорогенез изучали как в контроле, так и в обработанном материале. Растения, обработанные ЭМС, проявляли интересную особенность частичной интер-мейоцитной миграции хроматина при формировании каналов, образовании клювообразного выступа или прямого слияния клеток. Другим обнаруженным в исследовании явлением было слияние тетрад из-за растворения стенки. Феномен цитомиксиса отмечался почти на всех стадиях микроспорогенеза и затрагивал от нескольких до многих мейоцитов. Другие аномалии, такие как отставания, преждевременные движения, мости и неразделения хромосом, отмечались с незначительной частотой. Феномен цитомиксиса возрастал с увеличением длительности обработки ЭМС. Клетки с этими типами цитомиктических нарушений могут, вероятно, приводить к нерегулярному формированию гамет или зигот, гетерогенных по размеру пыльцевых зерен, или даже к потере фертильности в будущем.

P. Srivastava, G. Kumar

ЦИТОМІКТИЧНА ВАРИАБЕЛЬНІСТЬ
У САФЛОРА (*CARTHAMUS TINCTORIUS* L.),
ІНДУКОВАНОГО ЕМС

Насіння сафлору (*Carthamus tinctorius* L.) обробляли 0.5%-ним етилметансульфонатом (ЕМС) в одному з трьох режимів – 3, 5 і 7 год. Микроспорогенез вивчали як у контролі, так і в обробленому матеріалі. Рослини, оброблені ЕМС, виявляли цікаву особливість часткової інтер-мейоцитної міграції хроматину при формуванні каналів, утворенні дзюбоподібного чи прямому злитті клітин. Іншим виявленим в дослідженні явищем було злиття тетрад через розчинення стінки. Феномен цитоміксиса відзначався майже на всіх стадіях микроспорогенезу і зачіпав від кількох до багатьох мейоцитів. Інші аномалії, такі як відставання, передчасні рухи, мости і нерозділення хромосом, зустрічались з незначною частотою. Феномен цитоміксису зростав із збільшенням тривалості обробки ЕМС, Клітки з цими типами цитоміктичних порушень можуть, ймовірно, приводити до нерегулярного формування гамет або зигот, гетерогенних за розміром пилоквих зерен, або навіть до втрати фертильності в майбутньому.

REFERENCES

- Kornicke M. *Über ortsveränderung von Zellkernern* S BNiederhein, Ges Natur-U Heilkunde Bonn A, 1901, pp. 14–25.
- Gottschalk W., 1970. Chromosome and nucleus migration during microsporogenesis of *Pisum sativum*. *The Nucleus* **13**:1–9.
- Cheng K.C., Nieh H.W., Yang C.L., Wang I.M., Chou I.S., Chen J.S. 1975. Light and electron microscopical observations on cytomixis and the study of its relation to evolution; *Acta. Bot. Sin.* **17**: 60–69.
- Omara M.K., 1976. Cytomixis in *Lolium perenne*. *Chromosoma* **55**:267–271.
- Guochang Z., Quinglan Y. and Yongren Z., 1987. The relationship between cytomixis, chromosome mutation and karyotype evolution in Lily. *Caryologia* **40**: 243–259.
- Bedi Y.S., 1990. Cytomixis in woody species; *Proc. Indian Acad. Sci. (Plant Sci.)*. **100**: 233–238.
- Bellucci M., Roscini C. and Mariani A., 2003. Cytomixis in pollen mother cells of *Medicago sativa* L. *J. Hered.* **94**:512–516.
- Kaul M.L.H. and Nirmala C., 1991. Male sterile gene action diversity in barley and pea. *Nucleus* **34**: 32–39.
- Kamra O.P., 1960. Chromatin extrusion and cytomixis in pollen mother cells of *Hordeum*. *Hereditas* **46**: 562.
- Sinha A.R.P., 1988. Morphological and cytological changes induced in *Lindenbergia indica* following treatments. *Genet. Polonica* **29**:335–339.
- Amer S.M. and Mikhael E., 1986. Cytological effects of pesticides. XVI. Effect of the insecticide rotenone on root-mitosis of *Vicia faba*. *Cytologia* **51**: 171–176.
- Koul K.K., 1990. Cytomixis in pollen mother cells of *Alopecurus arundinaceus* Poir. *Cytologia* **55**: 169–173.
- Amma C.K.S., Namboodiri A.N., Panikkar A.O.N. and Sethuraj M.R., 1990. Radiation induced male sterility in *Hevea brasiliensis* (Willd exAdr. De Juss) Muell Agr. *Cytologia* **55**: 547–551.
- Nirmala C. and Kaul M., 1994. Male sterility in Pea IV. *Cytologia* **59**: 195–201.
- Haroun S.A., Al Shehri A.M. and Al Wadie H.M., 2004. Cytomixis in the microsporogenesis of *Vicia faba* L. (Fabaceae). *Cytologia* **69**: 7–11.
- Srivastav P.K. and Raina S.N., 1980. Cytomixis in *Clitoria termatea* L. var. *Pleniflora* Fantz. f. *Pleniflora*. *Curr. Sci.* **49**: 824–835.
- Zheng G.C., 1983. Cytomixis and chromosomal aberrations in the light of plant evolution; in *Select topic of evolution theory* (eds) S.C. Cheng, Z.Z. Huang (Beijing: Science Press) pp 60–67.
- de Nittancourt D. and Grant W.F., 1964. La cytogenetique de *Lotus (Leguminosae)* III. Un cas de cytomixie dans un hybride interspecific que; *Cytologia* **29**:191–195.
- Salesses G., 1970. Sur La phenomeane de cytomixie cher des hybrids triploides de prainier. Consequences genetiquez possibles. *Ann. Amelior Plant.* **20**:383–388.
- Mantu D.E. and Sharma A.K., 1983. Cytomixis in pollen mother cells of an apomictic ornamental *Ervatamia diraricata* Linn., Alston. *Cytologia* **48**:201–207.
- Sheida I M., Riahi H. and Hakim, M., 1993: Cytomixis in *Asparagus* L. *The Nucleus* **36**: 59–62.
- Diaz Lifante Z., Luque T., Santa Barbara C., 1992. Chromosome numbers of plants collected during Iter Meditranium II in Israel. *Bocconeia* **3**:229–250.
- Sheidai M., Arman M. and Zehzad B., 2002. Chromosome pairing and B-chromosomes in some *Aegilops* (Poaceae) species and populations from Iran. *Caryologia* **55**: 361–271.
- Sheidai M., Sheidai A., Masoomeh K.R., 2006. Cytology of some Iranian *Stipa* (Poaceae) species and populations. *Acta Bot. Croat.* **65** (1): 1–11.
- Consolaro M.E.L. and Pagliarini M.S. 1995. Cytomixis in pollen mother cells of *Centella asiatica* L. urban. *Nucleus* **38**:80–85.
- Veilleux R. 1985. Diploid and polyploid gametes in crop plants: Mechanism of formation and utilization in plant breeding. *Plant Breeding Rev.* **3**:253–288.
- Cheng K.C., Nie X.W., Wang Y.X. and Yang Q.L., 1980. The relation between cytomixis and variation of chromosome numbers in pollen mother cells of rye (*Secale cereals* L.). *Acta. Bot. Sin.* **22**: 216–220.
- Cheng K.C., Quiglan Y. and Yongsen Z., 1987. The relationship between cytomixis, chromosome mutation and karyotype evolution in Lily. *Caryologia* **40**:243–259
- Dagne K., 1994. Meiosis in interspecific hybrids and genomic interrelationships in *Guizotia* Cass. (Compositae). *Hereditas* **121**: 119–129.
- Bione N.C.P, Pagliarini M.S., De Toledo J.F.F., 2000. Meiotic behavior of several Brazilian soybean varieties. *Genet. Mol. Biol.* **23**(3): 623–631.
- Ghanima A.M. and Talaat A.A., 2003. Cytomixis and its possible evolutionary role in a Kuwait population of *Diplofaxisharra* (Boraginaceae). *Bot. J. Linn. Soc.* **143**: 169–175.
- Heslop-Harrison J., 1966. Cytoplasmic connexions between angiosperm meiocytes. *Ann. Bot.* **30**: 221–234.
- Risueño M.C., Giménez-Martin G., López-Saez J.F. and Garcia M.I.R., 1969. Connexions between meiocytes in plants. *Cytologia* **34**: 262–272.
- Lattoo S.K., Khan S., Bamotra S. and Dhar A.K., 2006. Cytomixis impairs meiosis and influences reproductive success in Chlorophytum comosum (Thunb) Jacq. – an additional strategy and possible implications. *Jour. of Biosci.* **31**: 629–637.
- Boldrini K.R. and Pagliarini M.S., 2006. Cell fusion and cytomixis during microsporogenesis in *Brachiaria humidicola* (Poaceae). *South African Journal of Botany* **72**(3): 478–481.

36. Falistocco E., Tosti T., Falcinelli M., 1995. Cytomixis in pollen mother cells of diploid *Dactylis*, one of the origins of 2n gametes. *J. Heredity* **86**: 448–453.
37. Ghaffari G.M., 2006. Occurrence of diploid and polyploid microspores in *Sorghum bicolor* (Poaceae) is the result of cytomixis. *Afr. J. Biotechnol.* **5**: 1450–1453.
38. Maheshwari P., 1950. An introduction to the embryology of angiosperms New York: Mc Graw Hill.
39. Kumar G. and Rai P., 2006. Induced cytotoxic variability in soybean (*Glycine max* L.). *the nucleus* **49(1,2)**: 53–57.
40. Pagliarini M.S. and Pereira M.A.S., 1992. Meiotic studies in *Pilocarpus pennatifolius* Lem. (Rutaceae). *Cytologia* **57**: 231–235.
41. Pagliarini M.S., Pissinatt M.B. and Silva N., 1993. Chromosomal behaviour and seed production in *Chlorophytum comosum* (Liliaceae). *Cytologia* **58**: 433–437.
42. Consolaro M.E.L., Pagliarini M.S. and Chaves L.J., 1996. Meiotic behavior, pollen fertility and seed production in Brazilian populations of *Centella asiatica* (L.) Urban (Umbelliferae). *Cytologia* **61**: 375–381.
43. Khazanehdari K.A. and Jones G.H., 1997. The causes and consequences of meiotic irregularity in the leek (*Allium ampeloprasum* spp. *Porrum*): implications for fertility, quality and uniformity. *Euphytica* **93**: 313–319.

Received 18.10.10